

# On parameter derivatives of the associated Legendre function of the first kind (with applications to the construction of the associated Legendre function of the second kind of integer degree and order)

Radosław Szmytkowski

Atomic Physics Division, Department of Atomic Physics and Luminescence,  
Faculty of Applied Physics and Mathematics, Gdańsk University of Technology,  
Narutowicza 11/12, PL 80-233 Gdańsk, Poland  
email: radek@mif.pg.gda.pl

October 23, 2009

## Abstract

A relationship between partial derivatives of the associated Legendre function of the first kind with respect to its degree,  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ , and to its order,  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$ , is established for  $m, n \in \mathbb{N}$ . This relationship is used to deduce four new closed-form representations of  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  from those found recently for  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$  by the present author [R. Szmytkowski, J. Math. Chem. 46 (2009) 231]. Several new expressions for the associated Legendre function of the second kind of integer degree and order,  $Q_n^m(z)$ , suitable for numerical purposes, are also derived.

**KEY WORDS:** Legendre functions; parameter derivatives; special functions

**MSC2010:** 33C45, 33C05

## 1 Introduction

Recently, an interest has arisen in the derivation of closed-form expressions for parameter derivatives of the associated Legendre functions [1–4]. In Refs. [3, 4], the present author has extensively studied such derivatives for the associated Legendre function of the first kind,  $P_\nu^\mu(z)$ , in the case when one of the parameters is a fixed integer. In particular, in Ref. [3], using finite-sum expressions for  $P_n^\mu(z)$ , with  $n \in \mathbb{N}$ , we have found the following two representations of  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$ , with  $m \in \mathbb{N}$ :

$$\begin{aligned} \left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m} &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \\ &+ (-)^m \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^{m-1} (-)^k \frac{(k+n)!(m-k-1)!}{k!(n-k)!} \left( \frac{z-1}{2} \right)^k \\ &+ \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} \frac{(k+n+m)!\psi(k+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z-1}{2} \right)^k \quad (0 \leq m \leq n) \end{aligned} \quad (1.1)$$

and

$$\begin{aligned}
\left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m} &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} + [\psi(n+m+1) + \psi(n-m+1)] P_n^m(z) \\
&\quad - (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)! \psi(k+m+1)}{k!(k+m)!(n-k)!} \left( \frac{z+1}{2} \right)^k \\
&\quad (0 \leq m \leq n). \tag{1.2}
\end{aligned}$$

In turn, in Ref. [4], using contour-integral representations of  $\partial P_\nu^m(z)/\partial \nu$ , we have arrived, among others, at the following three expressions for  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ , again with  $m, n \in \mathbb{N}$ :

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} - [\psi(n+1) + \psi(n-m+1)] P_n^m(z) \\
&\quad + \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} \frac{(k+n+m)! \psi(k+n+m+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z-1}{2} \right)^k \\
&\quad + \frac{(n+m)!}{(n-m)!} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^n \frac{(k+n)! \psi(k+n+1)}{k!(k+m)!(n-k)!} \left( \frac{z-1}{2} \right)^k \\
&\quad (0 \leq m \leq n), \tag{1.3}
\end{aligned}$$

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} + [\psi(n+1) - \psi(n-m+1)] P_n^m(z) \\
&\quad - (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z+1}{2} \right)^k \\
&\quad + (-)^{n+m} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (-)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\
&\quad \quad \times [\psi(k+n+m+1) - \psi(k+m+1)] \left( \frac{z+1}{2} \right)^k \\
&\quad + (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\quad \quad \times [\psi(k+n+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k \quad (0 \leq m \leq n) \tag{1.4}
\end{aligned}$$

and

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} + [\psi(n+m+1) - \psi(n+1)] P_n^m(z) \\
&\quad - (-)^{n+m} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^{m-1} \frac{(k+n)!(m-k-1)!}{k!(n-k)!} \left( \frac{z+1}{2} \right)^k \\
&\quad + (-)^{n+m} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (-)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\
&\quad \quad \times [\psi(k+n+m+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k \\
&\quad + (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\quad \quad \times [\psi(k+n+1) - \psi(k+m+1)] \left( \frac{z+1}{2} \right)^k \quad (0 \leq m \leq n). \tag{1.5}
\end{aligned}$$

In the above equations, and in what follows,  $\psi(\zeta)$  is the digamma function defined as

$$\psi(\zeta) = \frac{1}{\Gamma(\zeta)} \frac{d\Gamma(\zeta)}{d\zeta}. \quad (1.6)$$

In the present paper, we shall pursue further the subject of derivation of closed-form expressions for parameter derivatives of the associated Legendre function of the first kind. First, in Section 2, we shall show that there exists a simple relationship between the derivatives  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  and  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$ , both with  $m, n \in \mathbb{N}$ . Next, in Section 3, we shall use this relationship, in conjunction with the formulas (1.1) and (1.2), to derive four further representations of  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ , two of them involving sums of powers of  $(z+1)/2$  and the remaining two — sums of powers of  $(z-1)/2$ . Interestingly, each out of these four representations contains only *two* sums. Therefore, the two new expressions for  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  containing sums of powers of  $(z+1)/2$  appear to be markedly simpler than the previously derived representations (1.4) and (1.5), while the two new expressions involving sums of powers of  $(z-1)/2$  are of the same degree of complexity as the representation (1.3). In addition, as a by-product, we shall obtain in that section two useful finite-sum representations of  $[\psi(n+m+1) - \psi(n+1)]P_n^m(z)$ . In the final Section 4, we shall exploit the results of Section 3 to find some new representations of the associated Legendre function of the second kind of integer degree and order,  $Q_n^m(z)$ , suitable for use for numerical purposes in various parts of the complex  $z$ -plane.

Throughout the paper, we shall be adopting the standard convention according to which  $z \in \mathbb{C} \setminus [-1, 1]$ , with the phases restricted by

$$-\pi < \arg(z) < \pi, \quad -\pi < \arg(z \pm 1) < \pi \quad (1.7)$$

(this corresponds to drawing a cut in the  $z$ -plane along the real axis from  $z = -\infty$  to  $z = +1$ ), hence,

$$-z = e^{\mp i\pi} z \quad -z+1 = e^{\mp i\pi}(z-1) \quad -z-1 = e^{\mp i\pi}(z+1) \quad (\arg(z) \gtrless 0). \quad (1.8)$$

Also, it will be implicit that  $x \in [-1, 1]$ ,  $\mu, \nu \in \mathbb{C}$  and  $k, m, n \in \mathbb{N}$ . Finally, it will be understood that if the upper limit of a sum is less by unity than the lower one, then the sum vanishes identically. The associated Legendre functions of the first and the second kinds used in the paper are those of Hobson [5] (cf. also Refs. [6–8]).

Before proceeding to the matter, we emphasize that the results obtained in the present paper are interesting not only for their own (mathematical) sake. The derivatives  $\partial P_\nu^m(z)/\partial \nu$  and  $\partial P_\nu^m(x)/\partial \nu$  are met in solutions of some boundary value problems of theoretical acoustics, electromagnetism, heat conduction and other branches of theoretical physics and applied mathematics (for a list of references illustrating this statement, see Ref. [4]). Physical applications of the associated Legendre functions of the second kind of integer degree and order,  $Q_n^m(z)$  and  $Q_n^m(x)$ , (for which, recall, several new explicit expressions are derived in Section 4) are even more abundant [9, 10].

## 2 A relationship between $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ and $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$

The departure point for our considerations in this section is the following Rodrigues-type formula, due to Barnes [11], for the associated Legendre function of the first kind when the sum of its degree and its order is a non-negative integer:

$$P_\nu^{m-\nu}(z) = \frac{1}{2^\nu \Gamma(\nu+1)} (z^2-1)^{(m-\nu)/2} \frac{d^m(z^2-1)^\nu}{dz^m}. \quad (2.1)$$

In terms of the Jacobi polynomial

$$P_n^{(\alpha, \beta)}(z) = \frac{1}{2^n n!} (z-1)^{-\alpha} (z+1)^{-\beta} \frac{d^n}{dz^n} [(z-1)^{n+\alpha} (z+1)^{n+\beta}] \quad (\alpha, \beta \in \mathbb{C}), \quad (2.2)$$

Eq. (2.1) may be rewritten as

$$P_\nu^{m-\nu}(z) = \frac{m!}{\Gamma(\nu+1)} \left( \frac{z^2-1}{4} \right)^{(\nu-m)/2} P_m^{(\nu-m, \nu-m)}(z). \quad (2.3)$$

(With no doubt, the reader has immediately realized that the Jacobi polynomial appearing on the right-hand side of Eq. (2.3) is a multiple of the Gegenbauer polynomial  $C_m^{(\nu-m+1/2)}(z)$ . However, we shall not make any use of this fact here.) Differentiation of Eq. (2.3) with respect to  $\nu$ , followed by setting  $\nu = n$ , yields

$$\begin{aligned} \left. \frac{\partial P_\nu^{m-n}(z)}{\partial \nu} \right|_{\nu=n} - \left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m-n} &= \frac{1}{2} P_n^{m-n}(z) \ln \frac{z^2-1}{4} - \psi(n+1) P_n^{m-n}(z) \\ &+ \frac{m!}{n!} \left( \frac{z^2-1}{4} \right)^{(n-m)/2} \left. \frac{\partial P_m^{(\lambda, \lambda)}(z)}{\partial \lambda} \right|_{\lambda=n-m}. \end{aligned} \quad (2.4)$$

The replacement of  $m$  by  $n+m$  results in the relationship

$$\begin{aligned} \left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} - \left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m} &= \frac{1}{2} P_n^m(z) \ln \frac{z^2-1}{4} - \psi(n+1) P_n^m(z) \\ &+ \frac{(n+m)!}{n!} \left( \frac{z^2-1}{4} \right)^{-m/2} \left. \frac{\partial P_{n+m}^{(\lambda, \lambda)}(z)}{\partial \lambda} \right|_{\lambda=-m}. \end{aligned} \quad (2.5)$$

If in Eq. (2.5) one exploits the following two explicit representations of the Jacobi polynomial  $P_n^{(\alpha, \beta)}(z)$ :

$$P_n^{(\alpha, \beta)}(z) = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+\alpha+\beta+1)} \sum_{k=0}^n \frac{\Gamma(k+n+\alpha+\beta+1)}{k!(n-k)!\Gamma(k+\alpha+1)} \left( \frac{z-1}{2} \right)^k, \quad (2.6)$$

$$P_n^{(\alpha, \beta)}(z) = (-)^n \frac{\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)} \sum_{k=0}^n (-)^k \frac{\Gamma(k+n+\alpha+\beta+1)}{k!(n-k)!\Gamma(k+\beta+1)} \left( \frac{z+1}{2} \right)^k, \quad (2.7)$$

after making use of Eq. (2.3) and of the easily provable relation

$$\lim_{\lambda \rightarrow -m} \frac{\psi(k+\lambda+1)}{\Gamma(k+\lambda+1)} = (-)^{k+m} (m-k-1)! \quad (0 \leq k \leq m-1), \quad (2.8)$$

one obtains

$$\begin{aligned} \left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} - \left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m} &= \frac{1}{2} P_n^m(z) \ln \frac{z^2-1}{4} - 2\psi(n-m+1) P_n^m(z) \\ &- (-)^m \frac{(n+m)!}{(n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} (-)^k \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z-1}{2} \right)^k \\ &+ \frac{(n+m)!}{(n-m)!} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^n \frac{(k+n)!}{k!(k+m)!(n-k)!} \\ &\times [2\psi(k+n+1) - \psi(k+1)] \left( \frac{z-1}{2} \right)^k \quad (0 \leq m \leq n) \end{aligned} \quad (2.9)$$

and

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} - \left. \frac{\partial P_n^\mu(z)}{\partial \mu} \right|_{\mu=m} &= \frac{1}{2} P_n^m(z) \ln \frac{z^2-1}{4} - 2\psi(n-m+1) P_n^m(z) \\
&- (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z+1}{2} \right)^k \\
&+ (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\times [2\psi(k+n+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k \quad (0 \leq m \leq n), \tag{2.10}
\end{aligned}$$

respectively.

### 3 Some new representations of $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ with $0 \leq m \leq n$

In this section, we shall use the results of Section 2 to provide several expressions for the derivative  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ , which differ from these given in Eqs. (1.3)–(1.5).

The first from among these expressions for  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  follows if one plugs the representation (1.2) of  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$  into the relationship (2.10). The result

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} + [\psi(n+m+1) - \psi(n-m+1)] P_n^m(z) \\
&- (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z+1}{2} \right)^k \\
&+ (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\times [2\psi(k+n+1) - \psi(k+m+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k \quad (0 \leq m \leq n) \tag{3.1}
\end{aligned}$$

is seen to be much simpler than either of the representations (1.4) or (1.5). An infinite variety of other representations of  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ , involving sums of powers of  $(z+1)/2$ , may be obtained by taking linear combinations, with coefficients such that their sum is unity, of the expressions in Eqs. (1.4), (1.5) and (3.1). For instance, multiplying Eq. (3.1) by  $-1$  and adding to the sum of Eqs. (1.4) and (1.5) leads to another remarkably simple formula

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} \\
&- (-)^{n+m} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^{m-1} \frac{(k+n)!(m-k-1)!}{k!(n-k)!} \left( \frac{z+1}{2} \right)^k \\
&+ (-)^{n+m} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (-)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\
&\times [2\psi(k+n+m+1) - \psi(k+m+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k \\
&(0 \leq m \leq n). \tag{3.2}
\end{aligned}$$

It is seen that for  $m = 0$  both the representations (3.1) and (3.2) of  $[\partial P_\nu^m(z)/\partial\nu]_{\nu=n}$  reduce to the formula

$$\left. \frac{\partial P_\nu(z)}{\partial\nu} \right|_{\nu=n} = P_n(z) \ln \frac{z+1}{2} + 2 \sum_{k=0}^n (-)^{k+n} \frac{(k+n)!}{(k!)^2(n-k)!} [\psi(k+n+1) - \psi(k+1)] \left( \frac{z+1}{2} \right)^k, \quad (3.3)$$

found by the present author in Ref. [12, Section 5.2.7] (cf. also Ref. [13]).

From the above findings, one may deduce two interesting and, as we shall see in a moment, useful identities involving the function  $P_n^m(z)$ . If we equate the right-hand sides of Eqs. (1.4) and (3.1), this results in the first of these relations:

$$\begin{aligned} [\psi(n+m+1) - \psi(n+1)]P_n^m(z) &= (-)^{n+m} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (-)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\ &\quad \times [\psi(k+n+m+1) - \psi(k+m+1)] \left( \frac{z+1}{2} \right)^k \\ &- (-)^n \frac{(n+m)!}{(n-m)!} \left( \frac{z+1}{z-1} \right)^{m/2} \sum_{k=0}^n (-)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\ &\quad \times [\psi(k+n+1) - \psi(k+m+1)] \left( \frac{z+1}{2} \right)^k \\ &\quad (0 \leq m \leq n). \end{aligned} \quad (3.4)$$

Replacement of  $z$  by  $-z$  in the above equation, followed by the use of the well-known property

$$P_n^m(-z) = (-)^n P_n^m(z) \quad (0 \leq m \leq n) \quad (3.5)$$

and also of Eq. (1.8), leads to the second identity:

$$\begin{aligned} [\psi(n+m+1) - \psi(n+1)]P_n^m(z) &= \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\ &\quad \times [\psi(k+n+m+1) - \psi(k+m+1)] \left( \frac{z-1}{2} \right)^k \\ &- \frac{(n+m)!}{(n-m)!} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^n \frac{(k+n)!}{k!(k+m)!(n-k)!} \\ &\quad \times [\psi(k+n+1) - \psi(k+m+1)] \left( \frac{z-1}{2} \right)^k \\ &\quad (0 \leq m \leq n). \end{aligned} \quad (3.6)$$

Playing with Eq. (1.3) and with the identity (3.6), one may obtain an infinite variety of representations of  $[\partial P_\nu^m(z)/\partial\nu]_{\nu=n}$  containing sums of powers of  $(z-1)/2$ . Two examples of such representations are

$$\begin{aligned} \left. \frac{\partial P_\nu^m(z)}{\partial\nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} - [\psi(n+m+1) + \psi(n-m+1)]P_n^m(z) \\ &+ \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\ &\quad \times [2\psi(k+n+m+1) - \psi(k+m+1)] \left( \frac{z-1}{2} \right)^k \\ &+ \frac{(n+m)!}{(n-m)!} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^n \frac{(k+n)! \psi(k+m+1)}{k!(k+m)!(n-k)!} \left( \frac{z-1}{2} \right)^k \\ &\quad (0 \leq m \leq n) \end{aligned} \quad (3.7)$$

and

$$\begin{aligned}
\left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} &= P_n^m(z) \ln \frac{z+1}{2} + [\psi(n+m+1) - 2\psi(n+1) - \psi(n-m+1)] P_n^m(z) \\
&+ \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} \frac{(k+n+m)! \psi(k+m+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z-1}{2} \right)^k \\
&+ \frac{(n+m)!}{(n-m)!} \left( \frac{z-1}{z+1} \right)^{m/2} \sum_{k=0}^n \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\times [2\psi(k+n+1) - \psi(k+m+1)] \left( \frac{z-1}{2} \right)^k \quad (0 \leq m \leq n). \quad (3.8)
\end{aligned}$$

For  $m = 0$ , both Eqs. (3.7) and (3.8) reduce to the Schelkunoff's formula [14] (cf. also Ref. [12, Section 5.2.6])

$$\left. \frac{\partial P_\nu(z)}{\partial \nu} \right|_{\nu=n} = P_n(z) \ln \frac{z+1}{2} - 2\psi(n+1) P_n(z) + 2 \sum_{k=0}^n \frac{(k+n)! \psi(k+n+1)}{(k!)^2 (n-k)!} \left( \frac{z-1}{2} \right)^k. \quad (3.9)$$

From the representations of  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  found above, one may construct counterpart representations for  $[\partial P_\nu^{-m}(z)/\partial \nu]_{\nu=n}$ , using the relationship [4]

$$\left. \frac{\partial P_\nu^{-m}(z)}{\partial \nu} \right|_{\nu=n} = \frac{(n-m)!}{(n+m)!} \left. \frac{\partial P_\nu^m(z)}{\partial \nu} \right|_{\nu=n} - [\psi(n+m+1) - \psi(n-m+1)] P_n^{-m}(z) \quad (0 \leq m \leq n) \quad (3.10)$$

and the well-known property

$$P_n^m(z) = \frac{(n+m)!}{(n-m)!} P_n^{-m}(z) \quad (0 \leq m \leq n). \quad (3.11)$$

Moreover, it does not offer any difficulty to derive counterpart expressions on the cut  $x \in [-1, 1]$  by using the formulas

$$\begin{aligned}
\left. \frac{\partial P_\nu^{\pm m}(x)}{\partial \nu} \right|_{\nu=n} &= e^{\pm i\pi m/2} \left. \frac{\partial P_\nu^{\pm m}(x+i0)}{\partial \nu} \right|_{\nu=n} = e^{\mp i\pi m/2} \left. \frac{\partial P_\nu^{\pm m}(x-i0)}{\partial \nu} \right|_{\nu=n} \\
&= \frac{1}{2} \left[ e^{\pm i\pi m/2} \left. \frac{\partial P_\nu^{\pm m}(x+i0)}{\partial \nu} \right|_{\nu=n} + e^{\mp i\pi m/2} \left. \frac{\partial P_\nu^{\pm m}(x-i0)}{\partial \nu} \right|_{\nu=n} \right] \quad (3.12)
\end{aligned}$$

together with

$$x+1 \pm i0 = 1+x, \quad x-1 \pm i0 = e^{\pm i\pi} (1-x). \quad (3.13)$$

Concluding this section we note that, in principle, the relations (2.9) and (2.10) might be also used in the opposite direction, i.e., to construct representations for  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$  from those known for  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$ . As it appears, however, that all expressions for  $[\partial P_n^\mu(z)/\partial \mu]_{\mu=m}$  obtainable in this way are much more complex (and thus potentially less useful) than these in Eqs. (1.1) and (1.2), we do not present them here.

## 4 Some new representations of $Q_n^m(z)$ with $0 \leq m \leq n$

The associated Legendre function of the second kind,  $Q_\nu^\mu(z)$ , may be defined [5] as the following linear combination of the Legendre functions of the first kind  $P_\nu^\mu(z)$  and  $P_\nu^\mu(-z)$ :

$$Q_\nu^\mu(z) = \frac{\pi}{2} e^{i\pi\mu} \frac{e^{\mp i\pi\nu} P_\nu^\mu(z) - P_\nu^\mu(-z)}{\sin[\pi(\nu+\mu)]} \quad (\text{Im}(z) \geq 0). \quad (4.1)$$

In the special case of  $\mu = m$ , Eq. (4.1) simplifies to

$$Q_\nu^m(z) = \frac{\pi e^{\mp i\pi\nu} P_\nu^m(z) - P_\nu^m(-z)}{2 \sin(\pi\nu)} \quad (\text{Im}(z) \geq 0), \quad (4.2)$$

hence, after exploiting the l'Hospital rule, one obtains

$$Q_n^m(z) = \mp \frac{1}{2} i\pi P_n^m(z) + \frac{1}{2} \frac{\partial P_\nu^m(z)}{\partial \nu} \Big|_{\nu=n} - \frac{(-)^n}{2} \frac{\partial P_\nu^m(-z)}{\partial \nu} \Big|_{\nu=n} \quad (0 \leq m \leq n, \text{Im}(z) \geq 0). \quad (4.3)$$

Thus, we see that the problem of evaluation of  $Q_n^m(z)$  with  $0 \leq m \leq n$  may be reduced to that of derivation of expressions for  $[\partial P_\nu^m(\pm z)/\partial \nu]_{\nu=n}$ .

Accordingly, after combining Eq. (4.3) with Eqs. (1.3) and (3.1), we obtain

$$\begin{aligned} Q_n^m(z) &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \mp \frac{1}{2} [\psi(n+m+1) + \psi(n+1)] P_n^m(z) \\ &\pm \frac{(\pm)^n (\mp)^m (n+m)!}{2 (n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} (\mp)^k \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\ &\pm \frac{(\pm)^{n+m}}{2} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (\pm)^k \frac{(k+n+m)! \psi(k+n+m+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\ &\mp \frac{(\pm)^n (n+m)!}{2 (n-m)!} \left( \frac{z \mp 1}{z \pm 1} \right)^{m/2} \sum_{k=0}^n (\pm)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\ &\quad \times [\psi(k+n+1) - \psi(k+m+1) - \psi(k+1)] \left( \frac{z \mp 1}{2} \right)^k \quad (0 \leq m \leq n), \end{aligned} \quad (4.4)$$

where the upper signs follow if  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  is evaluated from Eq. (1.3) and  $[\partial P_\nu^m(-z)/\partial \nu]_{\nu=n}$  from Eq. (3.1), while the lower signs result if the roles of Eqs. (1.3) and (3.1) are interchanged. The same expression for  $Q_n^m(z)$  as above is obtained if Eq. (4.3) is coupled with Eqs. (1.4) and (3.7). Further, using Eqs. (1.3) and (3.2) in Eq. (4.3) leads to

$$\begin{aligned} Q_n^m(z) &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \mp \frac{1}{2} [\psi(n+1) + \psi(n-m+1)] P_n^m(z) \\ &\pm \frac{(\pm)^n (-)^m}{2} \left( \frac{z \pm 1}{z \mp 1} \right)^{m/2} \sum_{k=0}^{m-1} (\mp)^k \frac{(k+n)!(m-k-1)!}{k!(n-k)!} \left( \frac{z \mp 1}{2} \right)^k \\ &\mp \frac{(\pm)^{n+m}}{2} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (\pm)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\ &\quad \times [\psi(k+n+m+1) - \psi(k+m+1) - \psi(k+1)] \left( \frac{z \mp 1}{2} \right)^k \\ &\pm \frac{(\pm)^n (n+m)!}{2 (n-m)!} \left( \frac{z \mp 1}{z \pm 1} \right)^{m/2} \sum_{k=0}^n (\pm)^k \frac{(k+n)! \psi(k+n+1)}{k!(k+m)!(n-k)!} \left( \frac{z \mp 1}{2} \right)^k \\ &\quad (0 \leq m \leq n). \end{aligned} \quad (4.5)$$

Next, if  $[\partial P_\nu^m(z)/\partial \nu]_{\nu=n}$  is obtained from Eq. (3.1) and  $[\partial P_\nu^m(-z)/\partial \nu]_{\nu=n}$  from Eq. (3.7), or vice

versa, then Eq. (4.3) yields the expressions

$$\begin{aligned}
Q_n^m(z) &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \mp \psi(n+m+1) P_n^m(z) \\
&\pm \frac{(\pm)^n (\mp)^m (n+m)!}{2 (n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} (\mp)^k \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&\pm \frac{(\pm)^{n+m}}{2} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (\pm)^k \frac{(k+n+m)!}{k!(k+m)!(n-m-k)!} \\
&\quad \times [2\psi(k+n+m+1) - \psi(k+m+1)] \left( \frac{z \mp 1}{2} \right)^k \\
&\mp \frac{(\pm)^n (n+m)!}{2 (n-m)!} \left( \frac{z \mp 1}{z \pm 1} \right)^{m/2} \sum_{k=0}^n (\pm)^k \frac{(k+n)!}{k!(k+m)!(n-k)!} \\
&\quad \times [2\psi(k+n+1) - 2\psi(k+m+1) - \psi(k+1)] \left( \frac{z \mp 1}{2} \right)^k \quad (0 \leq m \leq n). \quad (4.6)
\end{aligned}$$

We are not aware of any appearance of either of the formulas (4.4)–(4.6) in the literature. Furthermore, if Eqs. (3.1) and (3.8) are used in Eq. (4.3), this results in

$$\begin{aligned}
Q_n^m(z) &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \mp \psi(n+1) P_n^m(z) \\
&\pm \frac{(\pm)^n (\mp)^m (n+m)!}{2 (n-m)!} \left( \frac{z^2-1}{4} \right)^{-m/2} \sum_{k=0}^{m-1} (\mp)^k \frac{(k+n-m)!(m-k-1)!}{k!(n+m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&\pm \frac{(\pm)^{n+m}}{2} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (\pm)^k \frac{(k+n+m)! \psi(k+m+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&\pm \frac{(\pm)^n (n+m)!}{2 (n-m)!} \left( \frac{z \mp 1}{z \pm 1} \right)^{m/2} \sum_{k=0}^n (\pm)^k \frac{(k+n)! \psi(k+1)}{k!(k+m)!(n-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&(0 \leq m \leq n), \quad (4.7)
\end{aligned}$$

which is the same what follows if Eqs. (1.3) and (1.4) are plugged into Eq. (4.3) (cf. Ref. [4]). Finally, insertion of Eqs. (3.2) and (3.7) into Eq. (4.3) leads to

$$\begin{aligned}
Q_n^m(z) &= \frac{1}{2} P_n^m(z) \ln \frac{z+1}{z-1} \mp \frac{1}{2} [\psi(n+m+1) + \psi(n-m+1)] P_n^m(z) \\
&\pm \frac{(\pm)^n (-)^m}{2} \left( \frac{z \pm 1}{z \mp 1} \right)^{m/2} \sum_{k=0}^{m-1} (\mp)^k \frac{(k+n)!(m-k-1)!}{k!(n-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&\pm \frac{(\pm)^{n+m}}{2} \left( \frac{z^2-1}{4} \right)^{m/2} \sum_{k=0}^{n-m} (\pm)^k \frac{(k+n+m)! \psi(k+1)}{k!(k+m)!(n-m-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&\pm \frac{(\pm)^n (n+m)!}{2 (n-m)!} \left( \frac{z \mp 1}{z \pm 1} \right)^{m/2} \sum_{k=0}^n (\pm)^k \frac{(k+n)! \psi(k+m+1)}{k!(k+m)!(n-k)!} \left( \frac{z \mp 1}{2} \right)^k \\
&(0 \leq m \leq n), \quad (4.8)
\end{aligned}$$

which, in turn, is the same what is obtained if Eqs. (1.3) and (1.5) are coupled with Eq. (4.3) (cf. again Ref. [4]; for alternative derivations of the above result see Ref. [7, pages 81, 82 and 85] and Ref. [3]). Other expressions for  $Q_n^m(z)$  may be obtained by combining Eqs. (4.4)–(4.8), with the possible help of the identities (3.4) and (3.6).

Once the function  $Q_n^m(z)$  is known, one may find the function  $Q_n^{-m}(z)$  from the well-known relationship

$$Q_n^{-m}(z) = \frac{(n-m)!}{(n+m)!} Q_n^m(z) \quad (0 \leq m \leq n). \quad (4.9)$$

Using the formula

$$Q_n^{\pm m}(x) = \frac{(-)^m}{2} \left[ e^{\mp im/2} Q_n^{\pm m}(x + i0) + e^{\pm im/2} Q_n^{\pm m}(x - i0) \right], \quad (4.10)$$

which follows from the Hobson's [5] definition of the associated Legendre function of the second kind on the cut  $x \in [-1, 1]$ , and the identities in Eq. (3.13), from Eqs. (4.4)–(4.9) one may derive counterpart representations of  $Q_n^{\pm m}(x)$  with  $0 \leq m \leq n$ . Since the procedure does not offer any difficulty, we do not present the resulting expressions here.

## References

- [1] Yu. A. Brychkov, On the derivatives of the Legendre functions  $P_\nu^\mu(z)$  and  $Q_\nu^\mu$  with respect to  $\mu$  and  $\nu$ , *Integral Transforms Spec. Funct.* doi:10.1080/10652460903069660
- [2] H. S. Cohl, Derivatives with respect to the degree and order of associated Legendre functions for  $|z| > 1$  using modified Bessel functions, *Integral Transforms Spec. Funct.* in press
- [3] R. Szmytkowski, On the derivative of the associated Legendre function of the first kind of integer degree with respect to its order (with applications to the construction of the associated Legendre function of the second kind of integer degree and order), *J. Math. Chem.* 46 (2009) 231
- [4] R. Szmytkowski, On the derivative of the associated Legendre function of the first kind of integer order with respect to its degree (with applications to the construction of the associated Legendre function of the second kind of integer degree and order), preprint arXiv:0907.3217
- [5] E. W. Hobson, *The Theory of Spherical and Ellipsoidal Harmonics*, Cambridge University Press, Cambridge, 1931 [reprinted: (Chelsea, New York, 1955)]
- [6] L. Robin, *Fonctions Sphériques de Legendre et Fonctions Sphéroïdales*, vol. 1, Gauthier-Villars, Paris, 1957
- [7] L. Robin, *Fonctions Sphériques de Legendre et Fonctions Sphéroïdales*, vol. 2, Gauthier-Villars, Paris, 1958
- [8] L. Robin, *Fonctions Sphériques de Legendre et Fonctions Sphéroïdales*, vol. 3, Gauthier-Villars, Paris, 1959
- [9] P. M. Morse, H. Feshbach, *Methods of Theoretical Physics*, McGraw-Hill, New York, 1953
- [10] P. Moon, D. E. Spencer, *Field Theory for Engineers*, Van Nostrand, Princeton, 1961
- [11] E. W. Barnes, On generalized Legendre functions, *Quart. J. Pure Appl. Math.* 39 (1907) 97
- [12] R. Szmytkowski, On the derivative of the Legendre function of the first kind with respect to its degree, *J. Phys. A* 39 (2006) 15147 [erratum: 40 (2007) 7819]
- [13] R. Szmytkowski, Addendum to ‘On the derivative of the Legendre function of the first kind with respect to its degree,’ *J. Phys. A* 40 (2007) 14887. We note parenthetically that the simplest way to arrive at Eq. (5) in that paper is to differentiate both sides of the identity  $d^k z^\alpha / dz^k = [\Gamma(\alpha + 1) / \Gamma(\alpha - k + 1)] z^{\alpha - k}$  with respect to  $\alpha$ .
- [14] S. A. Schelkunoff, Theory of antennas of arbitrary size and shape, *Proc. IRE* 29 (1941) 493 [reprinted: *Proc. IEEE* 72 (1984) 1165]. Notice that Schelkunoff used a definition of the digamma function different from that adopted in Ref. [12] and in the present paper.